

# Game Theory Reading Notes

Eric Hsienchen Chu

May 11, 2023

This Reading Note is a part of [ECON699: Directed Study](#), Spring 2023. The reading material and contexts are greatly from Fudenberg and Tirole (1991), one of the most classic game theory textbooks. The document environment settings are generously provided by [Rebekah Dix](#), and I also use the 'xgames' package developed by [Prof. Ben Bernard](#) for game trees. I thank [Prof. Matteo Camboni](#) for his generous guidance of econ theory research and learning. All reports of typos/errors are welcome.

## Contents

<b>1</b>	<b>Static Games of Complete Information: Strategic-Form</b>	<b>3</b>
1.1	Introduction to Strategic-Form Games . . . . .	3
1.2	Nash Equilibrium . . . . .	3
1.3	Existence of Nash Equilibria . . . . .	4
<b>2</b>	<b>Iterated Strict Dominance, Rationalizability, and Correlated Equilibrium</b>	<b>5</b>
2.1	IESDS, Rationalizability . . . . .	5
2.2	Correlated Equilibrium . . . . .	6
2.3	Rationalizability and Subjective Correlated Equilibria . . . . .	6
<b>3</b>	<b>Dynamic Games of Complete Information: Extensive-Form</b>	<b>7</b>
3.1	Commitment and Perfection in Multi-Stage Games: Observable Action . . .	7
3.2	Extensive Form . . . . .	8
3.3	Strategies and Equilibria in Extensive-Form Games . . . . .	9
3.4	Backward Induction and Subgame Perfection . . . . .	10
<b>4</b>	<b>Applications of Multi-Stage Games with Observed Actions</b>	<b>11</b>
4.1	The Principle of Optimality and Subgame Perfection . . . . .	11
4.2	The Rubinstein-Stahl Bargaining Model . . . . .	12
4.3	Timing Games (Reading Examples) . . . . .	13
4.4	Iterated Conditional Dominance and Bargaining Game . . . . .	13
4.5	Open-loop and Closed-loop Equilibria . . . . .	13
<b>5</b>	<b>Repeated Games (skip)</b>	<b>14</b>

<b>6</b>	<b>Static Games of Incomplete Information: Bayesian Equilibrium</b>	<b>14</b>
6.1	Incomplete Information . . . . .	14
6.2	Public Good under Incomplete Information . . . . .	15
	<b>Appendix I. Introduction of Linear Programming</b>	<b>16</b>
	<b>Appendix II. Supermodular Games</b>	<b>17</b>

# 1 Static Games of Complete Information: Strategic-Form

## 1.1 Introduction to Strategic-Form Games

**Overview.** General framework:

- One-time Game: strategic-form games & equilibriums
- More than one period: extensive-form games & subgame perfect
- Uncertainty: incomplete information & dynamics

**Definition 1.1 (Static Game).** A static game  $\mathcal{G} = (\mathcal{I}, (S_i)_{i \in \mathcal{I}}, (u_i)_{i \in \mathcal{I}})$  has elements:

- (i) Finite set of Players:  $\mathcal{I} = \{1, \dots, n\}$
- (ii) A set  $S_i$  (pure strategy) available to Player  $i \forall i \in \mathcal{I}$
- (iii) Payoff function  $u_i : S \rightarrow \mathbb{R} \forall i \in \mathcal{I}$ , where  $S = S_1 \times \dots \times S_n$  is the set of pure strategy profile  $s = (s_1, \dots, s_n)$ .

**Remark.** Generally speaking, Action ( $\mathcal{A}_i$ ) should be different to Strategy ( $S_i$ ). **Only** in static games with complete information we can use  $\mathcal{A}_i = S_i$ .

Mixed Strategy:  $\sigma_i$

- probability distribution over  $S_i$ :  $\sigma_i(s_i) \leftarrow$  probability of Player  $i$  choosing strategy  $s_i$
  - Space of mixed strategy:  $\Sigma_i \Rightarrow \Sigma = \Sigma_1 \times \dots \times \Sigma_n$
- $\Rightarrow$  Player  $i$ 's payoff to mixed-profile  $\sigma$ :  $\sum_{s \in S} \left( \prod_{j=1}^{\mathcal{I}} \sigma_j(s_j) \right) u_i(s)$

**Definition 1.2 (Dominated Strategies).** Pure strategy  $S_i$  is strictly dominated for Player  $i$  if there exists  $\sigma'_i \in \Sigma_i$  such that  $u_i(\sigma'_i, s_{-i}) \succ u_i(S_i, s_{-i})$ .

**Remark.**  $S_i$  is dominated by any pure or mixed profile  $\sigma'_i$ .

## 1.2 Nash Equilibrium

**Definition 1.3.** A mixed-strategy profile  $\sigma^*$  is a NE if  $\forall$  player  $i$ , we have:  $u_i(\sigma_i^*, \sigma_{-i}^*) \geq u_i(s_i, \sigma_{-i}^*) \forall s_i \in S_i$ .

**Remark.**  $\sigma_i^*$  dominates all profile combination.

- NE is strict  $\Leftrightarrow \forall i$  and  $s_i \neq s_i^*$  (i.e.,  $s_i$  not optimal), we have  $u_i(s_i^*, s_{-i}^*) > u_i(s_i, s_{-i}^*)$ .
- NE is consistent.

### 1.3 Existence of Nash Equilibria

**Theorem 1.1 (Nash 1950b).** Every finite strategic-form game has a mixed-strategy equilibrium

**Remark.** Apply Kakutani's Fixed Point Theorem. Check:

- (i)  $\Sigma$  is a non-empty, convex, and compact (finite-dim) subset of Euclidean space.
- (ii) Reaction function  $r(\sigma)$  is non-empty and convex  $\forall \sigma$ .
- (iii)  $r$  has close graph: if  $(\sigma^n, \hat{\sigma}^n) \rightarrow (\sigma, \hat{\sigma})$  with  $\hat{\sigma}^n \in r(\sigma^n)$ , then  $\hat{\sigma} \in r(\sigma)$ .

Motivation: for  $\sigma$  (mixed-NE), players mutually respond best to the others. Say  $\Sigma$  is the mixed strategy profile, and that  $r : \Sigma \rightrightarrows \Sigma \equiv r_1(\sigma_{-1}) \times \cdots \times r_n(\sigma_{-n})$ . Then, a fixed point of  $r$  is a  $\sigma$  such that  $\sigma \in r(\sigma) \Rightarrow \sigma_i \in r_i(\sigma) \Leftrightarrow$  a fixed point of  $r$  is NE. From there, we can use "Kakutani's" to check.

- When payoff function is continuous  $\Rightarrow$  response correspondences have close graphs.
- An **upper semi-continuous** function has upward discontinuities.

**Theorem 1.2 (Pure NE- Debreu 1952; Glicksberg 1952; Fan 1952).** Consider a strategic-form game whose strategy space  $S_i$  are non-empty, compact, and convex subsets of an Euclidean space. If the payoff function  $u_i$  are continuous in  $s$  and **quasi-concave** in  $s_i$ , there exists a pure-strategy NE.

**Remark.** Payoff function is required to be continuous and quasi-concave here. If  $u_i$  conti, then it is non-empty by EVT (satisfying Cond 1) and has closed graph (satisfying Cond 3) instantaneously. If  $u_i$  is quasi-concave, it ensures a convex reaction correspondence.

**Theorem 1.3 (Mixed NE- Glicksberg 1952).** Consider a strategic-form game whose strategy space  $S_i$  are non-empty compact subsets of a metric space. If the payoff function  $u_i$  are continuous then there exists a mixed-strategy NE.

**Remark.** No compactness here ( $\Rightarrow$  not Euclidean space). No quasi-concave condition here. If  $u_i$  is disconti, then there is no Pure NE and may not have Mixed NE.

## 2 Iterated Strict Dominance, Rationalizability, and Correlated Equilibrium

### 2.1 IESDS, Rationalizability

**Motivation.** What strategies *could* a rational player play? Recall that in IESDS, we assume rational players will NEVER play those dominated (so we delete such strategies). Here, a rational agent play a **best response to some beliefs** about the opponents' strategies  $\Rightarrow$  **conjectures**

**Definition 2.1 (The Process of IESDS).** We can follow the steps:

- (i) Set  $S_i^0 \equiv S_i$  and  $\Sigma_i^0 \equiv \Sigma_i$ .
- (ii) Define  $S_i^n$  recursively by:  

$$S_i^n = \{s_i \in S_i^{n-1} \mid \text{exists no } \sigma_i \in \Sigma_i^{n-1} \text{ s.t. } u_i(\sigma_i, s_{-i}) > u_i(s_i, s_{-i}) \forall s_{-i} \in S_{-i}^{n-1}\}$$
- (iii) And, define  $\Sigma_i^n = \{\sigma_i \in \Sigma_i \mid \sigma_i(s_i) > 0 \text{ only if } s_i \in S_i^n\}$
- (iv) Set  $S_i^\infty = \bigcap_{n=0}^{\infty} S_i^n$  is the set of Player  $i$ 's Pure strategies that survive IESDS.
- (v) Set  $\Sigma_i^\infty = \{\sigma_i \in \Sigma_i \mid \text{no } \sigma_i' \text{ s.t. } u_i(\sigma_i', s_{-i}) > u_i(\sigma_i, s_{-i}) \forall s_{-i} \in S_{-i}^\infty\}$  is the set of Player  $i$ 's Mixed strategies that survive IESDS.

**Definition 2.2.** A game is solvable by IESDS if, for each player  $i$ ,  $S_i^\infty$  is a singleton (i.e., a one-element set).

**Remark.** When the IESDS yields a unique strategy profile, this strategy profile is necessarily a NE (think the prisoner's dilemma).

**Definition 2.3 (Rationalizable Strategies).** Set  $\tilde{\Sigma}_i^0 \equiv \Sigma_i$ , and for each  $i$  recursively define:

$$\tilde{\Sigma}_i^n = \left\{ \sigma_i \in \tilde{\Sigma}_i^{n-1} \mid \exists \sigma_{-i} \in \times_{j \neq i} \text{convex hull}(\tilde{\Sigma}_j^{n-1}) \text{ s.t. } u_i(\sigma_i, \sigma_{-i}) \geq u_i(\sigma_i', \sigma_{-i}) \forall \sigma_i' \in \tilde{\Sigma}_i^{n-1} \right\}.$$

Then, the rationalizable strategies for player  $i$  are  $R_i = \bigcap_{n=0}^{\infty} \tilde{\Sigma}_i^n$ .

**Remark.**  $\tilde{\Sigma}_i^{n-1}$  is the set of  $i$ 's surviving strategies that are best response to some strategies in  $\tilde{\Sigma}_{-i}^{n-1}$ , i.e., opponents' surviving strategies through (IESDS) round  $n - 1$ .

- The **convex hull** of a set  $\tilde{\Sigma}$  is the smallest convex set that contains it.
- In general, rationalizability will yield sharper predictions than IESDS.
- Every NE is rationalizable.

**Theorem 2.1 (Rationalizability and IESDS- Pearce 1984).** Rationalizability and IESDS coincide in two-player game. Specifically, if  $\mathcal{I} = 1, 2$  (two players), then  $R_i = \Sigma_i$  for  $i = 1, 2$  (set of rationalizability = set of mixed strategies surviving IESDS).

**Remark.** This theorem hints that:

- $R_i \subseteq \Sigma_i$  for all  $i$  (so that when  $i = 2$ ,  $R_i = \Sigma_i$ )
- Set of rationalizable strategies survive IESDS (so is a subset to it).

## 2.2 Correlated Equilibrium

**Motivation.** Aumann (1974) motivates such an idea of a **correlated equilibrium** achieved by some signaling devices/models. Suppose a correlating device  $(\Omega, (\mathcal{T}_{i \in \mathcal{I}}, p))$ :

**Definition 2.4 (Correlated Equilibrium A).** A correlated equilibrium  $\zeta$  relative to information structure  $(\Omega, (\mathcal{T}_{i \in \mathcal{I}}, p))$  is a NE in strategies that are adapted to this info structure.

**Remark.** That is, the vector  $(\zeta_1, \dots, \zeta_n)$  is a correlated equilibrium if, for every player  $i$  and adapted strategy  $\tilde{\zeta}_i$ ,

$$\sum_{\omega \in \Omega} p(\omega) u_i(\zeta_i(\omega), \zeta_{-i}(\omega)) \geq \sum_{\omega \in \Omega} p(\omega) u_i(\tilde{\zeta}_i(\omega), \zeta_{-i}(\omega))$$

We require that  $\zeta_i$  maximize player  $i$ 's ex-ante payoffs.

**Definition 2.5 (Correlated Equilibrium B).** A correlated equilibrium is any probability distribution  $p(\cdot)$  over the pure strategies  $S_1 \times \dots \times S_{\mathcal{I}}$  such that, for every player  $i$  and every function  $d_i(\cdot)$  that maps  $S_i$  to  $S_i$ , we have:

$$\sum_{s \in S} p(s) u_i(s_i, s_{-i}) \geq \sum_{s \in S} p(s) u_i(d_i(s_i), s_{-i})$$

**Remark.** It is equivalent to consider:  $\sum_{s_{-i} \in S_{-i}} p(s_{-i} | s_i) u_i(s_i, s_{-i}) \geq \sum_{s_{-i} \in S_{-i}} p(s_{-i} | s_i) u_i(s'_i, s_{-i})$

**Summary.** Both Pure and Mixed NEs are correlated equilibrium. Pure:  $p(\cdot)$  degenerate. Mixed:  $p(\cdot)$  can be joint distribution. The set of correlated equilibrium is convex and is at least as large as the convex hull of NE. For the existence problem, also use fixed point theorems to deal with.

## 2.3 Rationalizability and Subjective Correlated Equilibria

**Summary.** In matching pennies, players are allowed to have inconsistent strategic beliefs, by rationalizability. Meanwhile, subjective correlated equilibrium allows the belief to be *completely* arbitrary. It does not capture the restrictions implied by common knowledge and thus is less restrictive than rationalizability (check adapted strategy definition on Fudenberg and Tirole (1991) p.60).

### 3 Dynamic Games of Complete Information: Extensive-Form

**Overview.** Use extensive-form games to capture the dynamic structure. For instance, entry and entry deterrence in IO and time-consistency problem in Macro. Very essential example "Stackelberg leader" duopoly: one producer chooses action (i.e., output level) first and the other follows up. In this case, NE is defined as the strategy profile that neither players would gain from switching strategies. *Backward induction* is the key to solve these kinds of scenarios, alongside the idea of *subgame-perfect equilibrium*.

#### 3.1 Commitment and Perfection in Multi-Stage Games: Observable Action

**Summary.** Multi-stage game: (1) all players knew the action chosen in previous stage  $0, \dots, k-1$ , and (2) all players move simultaneously in each stage  $k$  (including the singleton of "do nothing"). Some notations:

- Players:  $i \in \mathcal{I}$
- Stages:  $0, \dots, k, k+1$
- Action set (choice): players simultaneously choose actions from  $\mathcal{A}_i(h^0)$
- Observable action profile (stage 0):  $a^0 \equiv (a_1^0, \dots, a_{\mathcal{I}}^0)$
- History: let  $h^0 = \emptyset$ , then history  $h^1$  at the beginning of stage 1 is just  $h^1 = a^0$
- Next stage:  $\mathcal{A}_i(h^1)$  then will be the action set for possible second-stage. Iteratively,  $h^{k+1}$  will be the history at the end of stage  $k$ /by start of stage  $k+1$ :  $h^{k+1} = (a^0, \dots, a^k)$ .
- Pure strategy: let  $H^k$  be the all stage- $k$  history and let  $\mathcal{A}_i(H^k) = \bigcup_{h^k \in H^k} \mathcal{A}_i(h^k)$ . A pure strategy for player  $i$  is a sequence  $\{s_i^k\}_{k=0}^K$

**Summary.** Backward induction can be applied to **any** finite game of perfect information (finite: finite stages & finite actions each stage).

**Definition 3.1 (Subgame-Perfect Equilibrium: SPE).** A strategy profile  $s$  of a multi-stage game with observed actions is a subgame-perfect equilibrium **if** for every  $h^k$ , the restriction  $s|_{h^k}$  to  $\mathcal{G}(h^k)$  is a NE of  $\mathcal{G}(h^k)$ .

**Example 3.1 (Two-stage game with Cournot).** Firm 1 and Firm 2 both have a const average cost of \$2 per unit. Firm 1 decides whether to invest in new technology with \$0 per unit but the technology itself costs  $f$ . Firm 2 observes Firm 1's investment and then propose  $q_2$  to compete in cournot format (against  $q_1$  by Firm 1). Suppose demand curve is given:  $p(q) = 14 - q$ . Firm 1's payoff will be:  $((14 - 2) - q_1 - q_2)q_1$  if not invest and  $((14 - 0) - q_1 - q_2)q_1 - f$  if invest. Firm 2's will be:  $((14 - 2) - q_1 - q_2)q_2$ .

*Answer.* Consider each cases and take FOC for each Firm for reaction functions:

- Firm 1 not invest:  $\frac{\partial((14-2)-q_1-q_2)q_1}{\partial q_1} = 12 - 2q_1 - q_2 = 0 \Rightarrow r_1(q_2) = 6 - \frac{q_2}{2}$ . By symmetry,  $r_2(q_1) = 6 - \frac{q_1}{2}$ . Solve this and get (4, 4) with both payoffs of 16.
- Firm 1 invest:  $r_1(q_2) = 7 - \frac{q_2}{2}$  and  $r_2(q_1) = 6 - \frac{q_1}{2}$ . Solve this and get  $(\frac{16}{3}, \frac{10}{3})$ . Firm 1's payoff will be  $\frac{256}{9} - f$ .

Thus, as long as the fixed cost  $f$  satisfies  $\frac{256}{9} - f > 16 \Rightarrow f < \frac{112}{9} \approx 12.44$ , Firm 1 will choose to invest.

**Motivation.** Players can sometimes gain benefit from making a binding **commitment**. With more than one players, commitments can be of value since such commitment would alter a sequence of actions the opponents may play. Typically, the optimal reaction function will be decreasing function of opponent's action/output. Commitment can also be achieved by "moving earlier" as it appears in Stackelberg.


**Example 3.2 (Odyssey-Commitment).** A general can burn the bridges as a commitment not to retreat (i.e., no way back) and Odysseus has himself lashed to the mast and order his sailors to plug their ears with wax as a commitment not to go to the Siren's island. Once they make these commitments, the cost of turning back is often modeled as  $\infty$ .

### 3.2 Extensive Form

**Definition 3.2 (Elements for Extensive Form).** The extensive-form game should contain:


- A finite set of player  $i \in \mathcal{I}$
- The order of moves and active players: "game tree"  $i : X \rightarrow \mathcal{I}$
- Payoff function for each player of the move:  $u_i : \mathcal{Z} \rightarrow \mathbb{R}$
- A set of actions:  $\mathcal{A}(h)$  at information set  $h$
- A partition of history  $H$ : what each player knows when making moves
- Probability distribution over exogenous events: "Nature"  $\mathcal{N}$

**Definition 3.3 (Perfect Information).** Information is *perfect* if nature has no moves AND each information set is a *singleton*. Else, imperfect information.



**Replication.** Figure 3.4 in Fudenberg and Tirole (1991)





**Replication.** Figure 3.6 in Fudenberg and Tirole (1991)

**Remark** (Signaling game). Note that the Nature moves first and choose "type" for player 1 (i.e., private information). For instance, in Fudenberg and Tirole (1991) Figure 3.6, Nature determines that Player 1 will be "Tough" type with probability  $\vartheta_T = 0.6$  and be "Weak" type with probability  $\vartheta_W = 0.4$ . Player 2 observes the action but not knowing Player 1's type, which forms this *signaling game* (imperfect information).

### 3.3 Strategies and Equilibria in Extensive-Form Games

**Definition 3.4** (Reduced Strategic Form). The *reduced strategic form* of an extensive-form game is obtained by identifying equivalent pure strategies. And, two pure strategies  $s_i$  and  $s'_i$  are *equivalent* if they lead to the same probability distribution over outcomes for all pure strategies of opponents.

**Definition 3.5** (Behavior Strategy). Let  $\Delta(\mathcal{A}(h_i))$  be the probability distribution on  $\mathcal{A}(h_i)$ . A *behavior strategy* for player  $i$ , denoted  $\sigma_i$ , is an element of the Cartesian product  $\prod_{h_i \in H_i} \Delta(\mathcal{A}(h_i))$ .

**Remark.** A behavior strategy specifies a probability distribution over actions at each  $h_i$ , and probability distribution at different information sets are independent.

**Theorem 3.1** (Kuhn 1953). In a game of perfect recall, mixed strategies are equivalent to the unique behavior strategies they generates.


**Theorem 3.2** (Zermelo 1913; Kuhn 1953). A finite (extensive-form) game of perfect information has a pure-strategy NE.

**Theorem 3.3** (Minimax Theorem; von Neumann 1928). Every two-player zero-sum game in which every player has a *finite* number of pure strategies has a value in mixed strategies. The value in mixed strategies of a two-player zero-sum game is given by  $v$ :

$$v := \max_{\sigma_I \in \Sigma_I} \min_{\sigma_{II} \in \Sigma_{II}} U(\sigma_I, \sigma_{II}) = \min_{\sigma_{II} \in \Sigma_{II}} \max_{\sigma_I \in \Sigma_I} U(\sigma_I, \sigma_{II})$$

### 3.4 Backward Induction and Subgame Perfection

**Motivation.** Though complex extensive-form game can be represented by strategic form one (and thus the concept of NE can be used), Selton (1965) argued that some NE are *more reasonable* than others.



**Replication.** Figure 3.14 in Fudenberg and Tirole (1991)

For instance, in extensive form game, the equilibrium is  $(D, L)$ , but, in strategic form,  $(D, L)$  and  $(U, R)$  are both NE. We say  $(U, R)$  is *not credible* as it relies on **empty threat** that Player 2 will only play R. We see Backward Induction yields the "correct" answer.

**Definition 3.6 (Subgame).** A proper subgame  $\mathcal{G}$  of an extensive-form game  $T$  consists of a *single* node and all its successors in  $T$ , with the property that if  $x' \in \mathcal{G}$  and  $x'' \in h(x')$ , then  $x'' \in \mathcal{G}$ . the information sets and payoffs of the subgame are inherited from the original game.

**Remark.** The requirements that the subgame begins with a single node and heritage info sets imply  $x$  in original game must be a singleton information set, i.e.,  $h(x) = \{x\}$ .

**Definition 3.7 (Subgame-Perfect Equilibrium; SPE).** A behavior-strategy profile  $\sigma$  of an extensive-form game is a *subgame-perfect equilibrium* if the restriction of  $\sigma|x$  to  $\mathcal{G}(x)$  is a NE for every proper subgame  $\mathcal{G}$ .

**Remark.** Every new stage can begin a proper subgame. Subgame perfection coincides with backward induction in finite games of perfect information.

**Summary.** Subgame perfection supposes not only the players expect NE in all subgames but also that ALL players expect the SAME equilibria.

## 4 Applications of Multi-Stage Games with Observed Actions

**Overview.** Include "open-loop" and "close-loop" equilibria, finite-horizon and infinite-horizon games, and some dynamic optimization. Consider mostly games with an infinite horizon as they are better models to many (real) situations. For instance, the bargaining game of Rubinstein (1982). Though those bargaining models are of infinite horizon, there exist some actions— accept the offer or exiting from the market— to end the game.

### 4.1 The Principle of Optimality and Subgame Perfection

**Motivation.** Check whether there are any history  $h^t$  where Player  $i$  can gain by *deviating* from the actions prescribed by  $s_i$  at  $h^t$  to verify a strategy profile of a multi-stage game with observed actions is subgame perfect.

**Theorem 4.1 (One-stage deviation Principle for finite-horizon).** In a finite multi-stage game with observed actions, strategy profile  $s$  is subgame perfect **if and only if** it satisfies the one-stage-deviation condition, meaning that NO Player  $i$  can gain by deviating from  $s$  in a single stage and still confronting to  $s$  thereafter.

**Remark.** It is equivalent to say: profile  $s$  is SP **if and only if** there exists NO Player  $i$  and NO strategy  $\hat{s}_i$  that agrees with  $s_i$  except at a single stage  $t$  and history  $h^t$ , and such that  $\hat{s}_i$  is a better response to  $s_{-i}$  than  $s_i$  conditional on  $h^t$  being reached.

**Definition 4.1 (Continuity at Infinity).** A game is continuous at infinity if, for each Player  $i$ , the utility function  $u_i$  satisfies:

$$\sup_{h, \tilde{h} \text{ s.t. } h^t = \tilde{h}^t} |u_i(h) - u_i(\tilde{h})| \rightarrow 0 \text{ as } t \rightarrow \infty.$$

**Remark.** Events in distant future are relatively unimportant. Continuity at infinity is satisfied if the overall payoff is a discounted present-value sum  $g_i^t(a^t)$  and per-period payoffs are uniformly bounded, i.e., there is a  $\mathcal{M}$  such that  $\max_{t, a^t} |g_i^t(a^t)| < \mathcal{M}$ .

**Definition 4.2 (One-stage deviation Principle for infinite-horizon).** In an infinite-horizon multi-stage game with observed actions that is **continuous at infinity**, profile  $s$  is SP **if and only if** there exists NO Player  $i$  and NO strategy  $\hat{s}_i$  that agrees with  $s_i$  except at a single stage  $t$  and history  $h^t$ , and such that  $\hat{s}_i$  is a better response to  $s_{-i}$  than  $s_i$  conditional on  $h^t$  being reached.

**Remark.** This Theorem shows if  $s$  satisfies one-stage-deviation condition, then it cannot be improved by any finite sequence of deviations in any subgame. It is also the essential idea of the Principle of Optimality for discounted dynamic programming (also see Blackwell (1965) and Appendix I for basics of Linear Programming).

## 4.2 The Rubinstein-Stahl Bargaining Model

**Motivation.** Two players must agree on how to share a pie of size 1. In period 0, 2, 4, ...:

- Player 1 proposes a sharing rule  $(x, 1 - x)$ , and Player 2 can accept or reject.
  - If Player 2 accepts, then game **ends**.
  - If Player 2 rejects, then proposes a sharing rule  $(x, 1 - x)$  in next period.
    - \* If Player 1 accepts, then game **ends**.
    - \* If Player 1 rejects, then proposes a sharing rule  $(x, 1 - x)$  in next period. ...

Consider discount factors  $\delta_1^t, \delta_2^t \in (0, 1)$ , then payoffs are  $(\delta_1^t x, \delta_2^t (1 - x))$  if sharing rule  $(x, 1 - x)$  is accepted at round  $t$ .

**Example 4.1** (Profile 1: A NE but not SP). One NE can be: *Player 1 always propose  $x = 1$  and refuse all smaller shares; Player 2 also always offers  $x = 1$  and accept any offers.* Yet, this will not be SP. Why? If Player 2 rejects Player 1's first offer and offer Player 1 a share  $x > \delta_1$ , then Player 1 should accepts since the best possible outcome if Player 1 rejects is to receive full pie (recall: Player 2 also proposes  $x = 1$ ) but only worth  $\delta_1$  next round.

**Example 4.2** (Profile 2: SPE). Consider the following: *Player  $i$  always demands a share  $\frac{1 - \delta_j}{1 - \delta_i \delta_j}$  when it is their turn. They accept any share  $\geq \frac{\delta_i(1 - \delta_j)}{1 - \delta_i \delta_j}$  and refuse any smaller share.* Why? Note that Player  $i$ 's demand of share is:

$$\frac{(1 - \delta_j)}{1 - \delta_i \delta_j} = \frac{(1 - \delta_i \delta_j) + \delta_i \delta_j - \delta_j}{1 - \delta_i \delta_j} = 1 - \underbrace{\delta_j \frac{(1 - \delta_j)}{1 - \delta_i \delta_j}}_{\text{Player 2 will accept}},$$

which is the highest share for Player  $i$  that is accepted by Player  $j$  (Player  $i$  cannot make any lower offer as it will be accepted by  $j$ ). Player  $i$  would also hurt if proposing a higher share, getting reject, and waiting till next round when Player  $j$  demands  $\frac{1 - \delta_i}{1 - \delta_i \delta_j}$ , i.e., Player  $i$  gets  $\left(1 - \frac{1 - \delta_i}{1 - \delta_i \delta_j}\right)$ :

$$\delta_i \left(1 - \frac{1 - \delta_i}{1 - \delta_i \delta_j}\right) = \delta_i^2 \frac{1 - \delta_j}{1 - \delta_i \delta_j} < \frac{1 - \delta_j}{1 - \delta_i \delta_j}$$

It is thus optimal for Player  $i$  to accept any offer that at least receives  $\frac{\delta_i(1 - \delta_j)}{1 - \delta_i \delta_j}$ .

**Remark.** This Rubinstein (1982) extends Stahl (1971) by considering infinite-horizon game. Two drawbacks for the finite-horizon version: ① solution for finite version depends on the length of game and who makes offer in last round, and ② it is more natural to continue the bargain if the offer at the last round is rejected. The uniqueness of the Infinite-Horizon Equilibrium matters here (Fudenberg and Tirole (1991), p.115).

**Remark.** Comparative statics: as  $\delta_i \rightarrow 1$ , fixed  $\delta_j$ , we notice Player  $i$  gets the whole pie. If  $\delta_i \rightarrow 0$ , Player  $j$  would NOT get the whole pie if  $\delta_j < 1$ . Why? Due to Player  $i$ 's first-move advantage, even a myopic Player  $i$  would get a positive share. If the discount factors  $\delta_i = \delta_j = \delta$ , then Player  $i$  would get a payoff  $v_1 = \frac{1-\delta}{1-\delta^2} = \frac{1}{1+\delta} > \frac{1}{2}$  as  $\delta \in (0, 1)$ .

### 4.3 Timing Games (Reading Examples)

- War of Attrition: fighting for prize with cost per fighting period Fudenberg and Tirole (1991) p.119-126
- Preemption Game: opposite of war of attrition. Illustrative: the decision of to build a new plant or to host innovation if market only allow one such addition.

### 4.4 Iterated Conditional Dominance and Bargaining Game

**Definition 4.3.** In a multi-stage game with observed actions, action  $a_i^t$  is *conditionally dominated at stage  $t$  given history  $h^t$*  if, in the subgame beginning at  $h^t$ , every strategy for Player  $i$  that assigns positive probability to  $a_i^t$  is strictly dominated.

**Remark.** Iterated Conditional Dominance means the process of deleting every conditionally dominated actions in every subgame.

**Theorem 4.2.** In a Finite- or Infinite-horizon game of perfect information, no subgame-perfect strategy profile is removed by iterated conditional dominance.

**Remark.** For a detailed walk-through of iterated conditional dominance with unique solution, see Fudenberg and Tirole (1991) p.130.

### 4.5 Open-loop and Closed-loop Equilibria

**Definition 4.4 (Closed and Open-loop).** The terms are used to distinguish information structures in multi-stage game. Multi-stage game with observed actions corresponds to the *closed-loop* info structure or feedback strategy. *Open-loop* strategy happens when players cannot observe the play of their opponents, as opposed to a closed-loop structure.

**Example 4.3 (Two-period).** We can use game with continuous actions to illustrate the use of Open-loop equilibria as benchmarks for measuring strategic effects. For Player:  $i = 1, 2$ ,

- Action: choose  $a_i \in \mathcal{A}_i$  and  $b_i \in \mathcal{B}_i$  simultaneously ( $\mathcal{A}, \mathcal{B}$  some interval on  $\mathbb{R}$ )
- Payoff:  $u_i$  assumed to be differentiable and concave
- Open-looped equilibrium:  $a_i^*$  maximizes  $u_i((a_i, a_{-i}^*), b^*)$  and  $b_i^*$  maximizes  $u_i(a^*, (b_i, b_{-i}^*))$  so that the Interior solution:  $\frac{\partial u_i}{\partial a_i} = 0 = \frac{\partial u_i}{\partial b_i}$ .

- Closed-looped equilibrium:  $\frac{\partial u_i}{\partial a_i} - \frac{\partial u_i}{\partial b_{-i}} \frac{\partial b_{-i}^*}{\partial a_i} = 0$  for Player  $i$  (Note: the **extra term** highlights Player  $i$ 's strategic incentive of altering  $a_i$  to affect  $b_{-i}$ ).

**Definition 4.5.** Profile  $\sigma^*$  is " $\epsilon$ -Nash equilibrium" if for all Player  $i$  and strategy  $\sigma_i$ , we have  $u_i(\sigma_i^*, \sigma_{-i}^*) \geq u_i(\sigma_i, \sigma_{-i}^*) - \epsilon$ . The profile is  $\epsilon$ -perfect equilibrium if no players can gain more than  $\epsilon$  by deviating in any subgame.


**Remark.** Sort of the vibe of " $\epsilon$ -good" argument for uniform continuity.  $\epsilon$ -Nash equilibrium can be regarded as how we relate the structures of large finite- and infinite-horizon games to smooth the finite-to-infinite horizon limit when utilities is **continuous at infinity**. (also see  $\epsilon$ -equilibrium).

## 5 Repeated Games (skip)

## 6 Static Games of Incomplete Information: Bayesian Equilibrium

### 6.1 Incomplete Information

**Overview.** When some players do NOT know payoffs of the others, we say the game has *incomplete information*. Harsanyi (1968) proposed the "Bayesian" contexts of understanding toward nature's moves for Player 1's "type/cost." All the players have the same *prior* beliefs about the probability distribution on nature's move. Then, Harsanyi's *Bayesian Nash Equilibrium* is precisely the the NE here of imperfect information games.



**Replication.** Figure 6.3 in Fudenberg and Tirole (1991)

**Example 6.1** (Figure 6.3). The optimal strategy for Player 2 is  $y = 1$  (i.e., entry), if  $x < \frac{1}{2(1-p_1)}$  and  $y = 0$  (i.e., no entry), if  $x > \frac{1}{2(1-p_1)}$ . The best response for Player 1 with "low-cost type" is  $x = 1$  (i.e., build) if  $y < \frac{1}{2}$ ,  $x = 0$  (i.e., no build) if  $y > \frac{1}{2}$ , and  $x \in [0, 1]$  if  $y = \frac{1}{2}$ .

## 6.2 Public Good under Incomplete Information

**Example 6.2** (Palfrey and Rosenthal 1989). Two player decide simultaneously whether to contribute to the public good with a 0 – 1 decision. Player  $i$ 's cost is  $c$ .

		Player 2	
		Contribute	Don't
Player 1	Contribute	$1 - c_1, 1 - c_2$	$1 - c_1, 1$
	Don't	$1, 1 - c_2$	$0, 0$

**Replication.** Figure 6.4 in Fudenberg and Tirole (1991)

Both player believe (common knowledge) that  $c_i$  is drawn independently from the same continuous and strictly increasing *cumulative function*  $P(\cdot) = [\underline{c}, \bar{c}]$  where  $c < 1 < \bar{c}$  ( $\Rightarrow P(\underline{c}) = 0$  and  $P(\bar{c}) = 1$ ). The cost  $c_i$  is thus Player's "type." A pure strategy here is a function  $s_i(c_i)$ ,  $s : [\underline{c}, \bar{c}] \rightarrow \{0, 1\}$ . The payoff is  $u_i(s_i, s_j, c_i) = \max(s_1, s_2) - c_i s_i$ .

(Last edited: 05/11/2023, Fudenberg and Tirole (1991) p.211)

# Appendix I. Introduction of Linear Programming

**Illustrative problem:** Let's consider:

$$\begin{aligned} \min_{x_1, x_2} x_1 + 3x_2 \quad \text{subject to} \quad & x_1 + x_2 \geq 2, \\ & x_2 \geq 1, \\ & x_1 - x_2 \geq 3 \end{aligned}$$

Let some helper coeffs  $p_1, p_2, p_3 \geq 0$ , we can rewrite our desired lower bound  $B$ :

$$p_1 \cdot \underbrace{(x_1 + x_2 \geq 2)}_{\text{CSTR. 1}} + p_2 \cdot \underbrace{(x_2 \geq 1)}_{\text{CSTR. 2}} + p_3 \cdot \underbrace{(x_1 - x_2 \geq 3)}_{\text{CSTR. 3}} = x_1 + 3x_2 \geq B$$

But by comparing the coefficients to objective function ( $x_1 + 3x_2$ ), we require:

$$\begin{aligned} x_1 : \quad & p_1 + p_3 = 1 \\ x_2 : \quad & p_1 + p_2 - p_3 = 3 \\ & p_1, p_2, p_3 \geq 0 \\ \rightarrow B = & 2p_1 + 1p_2 + 3p_3 \end{aligned}$$

Now we notice that our original minimization problem (**primal**) can be transformed into the following maximization problem (**dual**):

$$\begin{aligned} \max_{p_1, p_2, p_3} 2p_1 + p_2 + 3p_3 \quad \text{subject to} \quad & p_1 + p_3 = 1, \\ & p_1 + p_2 - p_3 = 3, \\ & p_1, p_2, p_3 \geq 0 \end{aligned}$$

We call this derived optimization problem the **dual LP problem**.

**Theorem (Dual Problem).** If we transform the dual into an equivalent minimization problem and then form its dual, we obtain a problem equivalent to the original problem. In mathematical words,

$$\min c'x \text{ subject to } Ax \geq b \Leftrightarrow \max p'b \text{ subject to } p'A = c', p \geq 0.$$

These two problems have the same values.

**Remark.** For non-negative vector  $p \geq 0$ , we have:  $Ax \geq b \Rightarrow (p'A)x \geq p'b$ . Therefore, to minimize (get lower bound) of objective function, it is equivalent to think maximizing  $p'b$  by the derived constraint  $p'A = c'$  matching the coefficient of  $x$ .

- Min problem of budget constraint given optimal utility  $\Leftrightarrow$  Max problem of utility function given budget constraint



## Appendix II. Supermodular Games

**Overview.** Developed by Topkis (1979), applied to economic problem by Vives (1990) and then Milgrom and Roberts (1990). The idea is that in some games best response correspondences are *increasing* so that players' strategies are "strategic complements." Supermodular games are well-behaved- have pure strategy NE. The upper and lower bounds of the NE sets and the rationalizable sets coincide. A supermodular game does not require convexity and differentiability assumptions, and it needs an **order structure on strategy space, weak continuity on payoff functions**, and that the marginal utility of each player's strategy is **monotonic**.

**Notations.** Suppose each player  $i$ 's strategy set  $S_i$  is a subset of a finite-dim Euclidean space  $\mathbb{R}^{m_i}$ . Then we define  $S \equiv \prod_{i=1}^I S_i$  is a subset of  $\mathbb{R}^m$ , where  $m \equiv \sum_{i=1}^I m_i$ . Let  $x$  and  $y$  denote two vectors in some Euclidean Space  $\mathbb{R}^k$ . Also, let  $x \geq y$  if  $x_k \geq y_k$  for all  $k = 1, \dots, K$ , and let  $x > y$  if  $x \geq y$  and there exists  $k$  such that  $x_k > y_k$ . The order  $\geq$  is only a partial order: if a vector dominates another in one component but is dominated in another component, the vectors cannot be compared. Next, we consider some lattice idea and define the "meet" as  $x \wedge y$  and "join" as  $x \vee y$ .

$$\begin{aligned} x \wedge y &\equiv (\min(x_1, y_1), \dots, \min(x_k, y_k)) \\ x \vee y &\equiv (\max(x_1, y_1), \dots, \max(x_k, y_k)) \end{aligned}$$

$S$  is a sublattice of  $\mathbb{R}^m$  if  $s \in S$  and  $\tilde{s} \in S$  implies that  $s \wedge \tilde{s} \in S$  and  $s \vee \tilde{s} \in S$

**Definition (Increasing difference).**  $u_i(s_i, s_{-i})$  has *increasing difference* in  $(s_i, s_{-i})$  **if**, for all  $(s_i, \tilde{s}_i) \in S_i^2$  and  $(s_{-i}, \tilde{s}_{-i}) \in S_{-i}^2$  s.t.  $s_i \geq \tilde{s}_i$  and  $s_{-i} \geq \tilde{s}_{-i}$ , we have:

$$u_i(s_i, s_{-i}) - u_i(\tilde{s}_i, s_{-i}) > u_i(s_i, \tilde{s}_{-i}) - u_i(\tilde{s}_i, \tilde{s}_{-i})$$

**Remark.** Change the  $\geq$  to  $>$  in conditions for *strictly increasing diff*. This definition says that an increase in the strategies of player  $i$ 's rivals (i.e., from playing  $\tilde{s}_{-i}$  to  $s_{-i}$ ) raises the desirability of playing high strategy for player  $i$  (i.e., from playing  $\tilde{s}_i$  to  $s_i$ ).

**Definition (Supermodular).**  $u_i(s_i, s_{-i})$  is *supermodular* in  $s_i$  **if** for each  $s_{-i}$  we have:

$$u_i(s_i \wedge \tilde{s}_i, s_{-i}) + u_i(s_i \vee \tilde{s}_i, s_{-i}) \geq u_i(s_i, s_{-i}) + u_i(\tilde{s}_i, s_{-i})$$

for all  $(s_i, \tilde{s}_i) \in S_i^2$ . Strictly supermodular if changing the sign from  $\geq$  to  $>$ .

**Definition (Supermodular Game).** A *supermodular game* is such that, for each  $i$ ,  $S_i$  is a sublattice of  $\mathbb{R}^{m_i}$ ,  $u_i$  shows increasing differences in  $(s_i, s_{-i})$ , and  $u_i$  is supermodular in  $s_i$ .

**Remark.** Suppose  $u_i$  is twice continuously differentiable, then  $u_i$  is supermodular **if and only if**  $\partial^2 u_i / \partial s_i \partial s_k \geq 0$ .

**Example.** Consider à la Diamond (1982) search model with the following payoff function:

$$u_i(s) = \alpha s_i \sum_{j \neq i} s_j - c(s_i) \begin{cases} s_i : \text{player } i\text{'s search intensity} \\ c(s_i) : \text{cost of search} \\ s_i \sum_{j \neq i} s_j : \text{probability of finding a trade partner} \\ \alpha : \text{standalone gain if a partner is found} \end{cases}$$

Since  $\partial^2 u_i / \partial s_i \partial s_j = \alpha > 0$ , this search (game) is supermodular.

**Theorem (Topkis 1979 & Vives 1990).** (a) (Topkis 1979) If, for each  $i$ ,  $S_i$  is compact and  $u_i$  is upper semi-continuous in  $s_i$  for each  $s_{-i}$ . Suppose the game is supermodular, then the Pure NE strategy is non-empty (i.e.,  $\neq \emptyset$ ) and possesses greatest/least equilibrium  $\bar{s}$  and  $\underline{s}$ . (b) (Vives 1990) if furthermore the game is strictly supermodular, the set of NE is a non-empty "complete" sublattice ("complete" means sup and inf of any subset  $\in$  that set).

**Theorem (Topkis 1979).** Consider a supermodular game with strictly increasing differences. If  $s_i \in r_i^*(s_{-i})$ ,  $\tilde{s}_i \in r_i^*(\tilde{s}_{-i})$  and that  $s_{-i} \geq \tilde{s}_{-i}$ , then  $s_i \geq \tilde{s}_i$ .

**Theorem (Milgrom and Roberts 1990).** Consider a supermodular game such that, for each  $i$ ,  $S_i$  is a complete sublattice and is bounded, and such that  $u_i$  is continuous and is bounded above. Then, the IESDS strategies yield a set of strategies in which the greatest and the least elements are NE:  $\bar{s}$  and  $\underline{s}$ .

(For more detailed walk-through and examples, check Fudenberg and Tirole (1991), p.496.)

## References

- Blackwell, D. (1965). Discounted Dynamic Programming. *The Annals of Mathematical Statistics*, 36(1), 226–235. <http://www.jstor.org/stable/2238089>
- Diamond, P. A. (1982). Aggregate Demand Management in Search Equilibrium. *Journal of Political Economy*, 90(5), 881–894. <http://www.jstor.org/stable/1837124>
- Fudenberg, D., & Tirole, J. (1991). Game Theory. *MIT Press Books, The MIT Press*, 1.
- Harsanyi, J. C. (1968). Games with Incomplete Information Played by "Bayesian" Players, I-III. Part III. The Basic Probability Distribution of the Game. *Management Science*, 14(7), 486–502. <http://www.jstor.org/stable/2628894>
- Milgrom, P., & Roberts, J. (1990). Rationalizability, Learning, and Equilibrium in Games with Strategic Complementarities (58th ed.). *Econometrica*, 6, 1255–1277. <https://doi.org/10.2307/2938316>
- Rubinstein, A. (1982). Perfect Equilibrium in a Bargaining Model. *Econometrica*, 50(1), 97–109. <http://www.jstor.org/stable/1912531>
- Vives, X. (1990). Nash Equilibrium with Strategic Complementarities. *Journal of Mathematical Economics*, 19(3), 305–321. <https://EconPapers.repec.org/RePEc:eee:mateco:v:19:y:1990:i:3:p:305-321>